
Postharvest Disease Management

- *Principles and Treatments* -

James E. Adaskaveg
Professor

Department of Plant Pathology
University of California, Riverside

Postharvest decay organisms




Fungi (eukaryotic organisms):

- Most important
- Mostly members of the Ascomycetes and Fungi imperfecti
- Propagation and dissemination by abundantly produced, mostly asexual spores
- Infection through wounds or sometimes through intact fruit surface.


Bacteria (prokaryotic organisms):

- Mostly pathogens of vegetables
- *Erwinia carotovora* is the most important postharvest pathogen causing a soft rot.
- Infections only through wounds.


Major postharvest decays of pome fruits



Alternaria decay of Asian pear caused by *Alternaria* sp.



Penicillium decay of Bosc pear caused by *Penicillium expansum*



Anthrachnose of apple caused by *Colletotrichum acutatum*

Postharvest decays of stone fruits

Brown rot (*Monilinia fructicola*)


Gray mold (*Botrytis cinerea*)

Rhizopus rot (*Rhizopus stolonifer*)

Sour rot (*Geotrichum candidum*)



Brown rot Infection through wounds and of healthy tissues




Gray mold

Infection through wounds and of senescent tissues



Sour rot



Infection through wounds of ripe fruit



Rhizopus rot


Infection through wounds

Postharvest decays of citrus





Green mold caused by *Penicillium digitatum* (most important on citrus)

Penicillium spp. are wound pathogens



Blue mold caused by *P. italicum* and green mold




Brown rot caused by *Phytophthora* spp. Infection through intact tissue.






Penicillium soilage

Major postharvest decays of citrus






Sour rot caused by *Geotrichum citri-aurantii*



Alternaria decay caused by *Alternaria* sp.



Tear stain and anthracnose caused by *Colletotrichum gloeosporioides*




Stem end rot caused by *Lasiodiplodia theobromae* (*B. rhodina*)

Postharvest decays of pomegranates and kiwifruit

Gray mold caused by
Botrytis cinerea



Infection through flower parts



Infection through cut stem ends at harvest




Major postharvest decays of tomato



Gray mold

Decay caused by
Botrytis cinerea




Rhizopus rot

R. stolonifer



Sour rot



Geotrichum candidum

Infection through wounds
of ripe fruit

Infection through wounds

Postharvest decay organisms

Penetration through wounds – Wound pathogens:

- Most common
- Only minor wounds required (micro-wounds).
- Wounds commonly occur before harvest (insect injuries, wind damage, etc.) or more frequently during and after harvest during handling, transport, packaging.
- **Goal in postharvest handling: Minimize fruit injuries.**

Penetration of intact fruit:

- Through surface of mature fruit.
- Quiescent infections that are established early during fruit growth but remain inactive until the fruit matures.
- Colonization of flower parts, invasion of maturing fruit



Infection by postharvest decay fungi



Conidiophore
and conidia
(asexual spores)
of *Botrytis
cinerea*



Spore germination: requires water,
oxygen, and sometimes nutrients




Host infection:

Penetration (through wounds or directly), inter- and intracellular growth. Enzymatic activities dissolve host cell walls and contents. Sometimes production of toxins that kill host cells.

- The Disease Triangle of Plant Pathology -

- A re-occurring interaction of
host, pathogen and environment -

Physiology,
optimum
harvest date



Conducive parameters during
storage, transportation, marketing

Identification, biology, ecology

Principles of Plant Disease Management

- Preventative (population)
 - Avoidance of the pathogen (Cultural practices)
 - Host resistance (Resistant varieties)
 - Exclusion (Quarantines and *Sorting/Grading*)*
 - Eradication (Eliminating or reducing inoculum - *Sanitation*)*
 - Protection/Prevention (Chemical or biological or physical treatments – *Cold temperature*)*
 - Curative (individual)
 - Therapy (Physical or chemical treatments)
-

* - Main postharvest practices for susceptible crop.

Preventative Practices - Eradication


Sanitation washes using oxidizing materials (chlorine, ozone, peroxide, etc.)

Micro-organisms in stem punctures, pits, injuries, natural cracks, or bruises, residual activity

No

Disinfestation of non-injured commodity surfaces and of micro-organisms in water

Yes



Comparison between postharvest sanitation and fungicide treatments

Treatment	Delivery System	Sources	Activity	Advantages	Dis-advantages
Chlorine	Water	Gas or liquid (Cl ₂ or NaOCL)	Fruit surface/In solution	Inexpensive, effective at low rates	Sensitive to pH and organic load; corrosive; reactive
Chlorine dioxide	Water	On-site generation	Fruit surface/In solution	Less sensitive to organic load	Initial cost of equipment; corrosive; training
Ozone	Water (low solubility)/ Air	On-site generation	In solution, but poor solubility; Air: anti-sporulation	Non-chlorine based, no disposal issues	Poor water solubility, initial cost of equipment; corrosive; training
Acidified hydrogen peroxide	Water	Liquid (H ₂ O ₂)	Fruit surface/In solution; some wound activity	Less sensitive to organic load and pH, no disposal issues	Conc. limits, cost, some sensitivity to Cl, pH, and organic load
Postharvest fungicide (e.g., Scholar)	Water	Dry or liquid Formulation	Wound protection	Highly effective	Residues; safety concerns; export tolerances (MRLs)

Chlorination in a hydrocooler (re-circulating)



Critical factors

Concentration
Contact time
pH
Organic load
Temperature

Chlorination on a brush bed (non-re-circulating)



Concentration
Contact time
pH

Strategies of postharvest decay control for protection, suppression, or eradication of decay

Altering the micro-environment


- Treatments with indirect effects on pathogen:

Change in pH

- Treatments with direct effects on pathogen:


Biocontrols: Competition, antibiosis, parasitism

Fungicides: Direct toxicity



Altering the host physiology and susceptibility

- Indirect effect on pathogen
- Plant growth regulators (PGRs)
 - Gibberellin (citrus)
 - 2,4-D (citrus)
 - Ethylene biosynthesis inhibitors?
- Effective against weak pathogens




Altering the micro-environment

Treatments with indirect effects on the pathogen:

- Change in pH
- Alkaline solutions of borax, sodium carbonate (soda ash), and sodium bicarbonate
- Accumulation of acid in potential infection sites, (e.g. SO_2)

Disease triangle




Treatments with direct effects on pathogen:

Biocontrols: Competition, antibiosis, parasitism


Fungicides: Direct toxicity

Borax, sodium carbonate (soda ash), and sodium bicarbonate

- *Change in pH*
 - Accumulation of alkali in potential infection sites on fruit surface
- *Germination of pathogen spores is inhibited (fungistatic action)*
 - Heated solutions are more toxic
- *Disadvantages*
 - Change in pH is gradually reversed by acid fruit juice
 - Fruit staining
 - Fruit dehydration
 - No residual activity



Usage of borax, sodium carbonate (soda ash), and sodium bicarbonate in postharvest treatments of lemons





Usage of borax, sodium carbonate (soda ash), and sodium bicarbonate in postharvest treatments of lemons

Treatment with heated soda ash



Water rinse after soda ash treatment



Altering the micro-environment

Treatments with indirect effects on the pathogen:


Change in pH

Alkaline solutions of borax, sodium carbonate, and sodium bicarbonate

Treatments with direct effects on pathogen:

Biocontrols: Competition, antibiosis, parasitism

Fungicides: Direct toxicity



Biocontrols:

Competition, antibiosis, parasitism

- Development is driven by safety concerns
- Activity from laboratory experiments is difficult to transfer into a commercial scale
- No activity against existing infections (infections that occur at harvest)
- Efficacy is generally inconsistent and never complete
- Previously, 2 products registered:
 - Aspire (no longer manufactured), see NEXY (*Candida oleophila*)
 - Bio-Save (*Pseudomonas syringae*), still in use

Bio-Save 10 LP

ACTIVE INGREDIENT:

Pseudomonas syringae Strain ESC-10 29.8%

INERT INGREDIENTS:

Total 70.2%

Note: Contains a minimum of 9×10^{10} colony forming units per gram of formulated product.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

Precaution at Use: Strictly for use as directed. Read and follow all directions on the label. Do not use if the label is torn, missing, or faded.

Hazards to Humans and Domestic Animals: Avoid contact with skin, eyes and clothing. When mixing wear protective eye wear (goggles, face shield or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before re-use.

Environmental Hazards: Do not contaminate water when disposing of equipment wash waters or rinsate.

STATEMENT OF PRACTICAL TREATMENT

If in eyes, flush with plenty of water. Get medical attention if irritation persists.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Storage:** Store only in original containers under refrigeration conditions. Avoid heat or warm temperatures during storage or transportation. Keep refrigerated until used. Store product separately from foods.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Put empty container in trash. Do not re-use empty container.

WARRANTY STATEMENT

EcoScience Produce Systems Corp. warrants that the use of this product will result in the control of the pest named on the label. The use of this product beyond the control of the manufacturer, no guarantee, express or implied, is made as to the effects of use, or the results to be obtained if not used in accordance with directions or established safe practices. The buyer must assume all responsibilities, including injury or damage, resulting from its failure to follow all such or in combination with other materials.

ECOSCIENCE PRODUCE SYSTEMS CORP.'S LIABILITY FOR ANY MALFUNCTION OR NON-FUNCTION OF THIS PRODUCT SHALL BE LIMITED TO THE ACTUAL COST OF REPLACEMENT OF THE PRODUCT, AND SHALL NOT, IN ANY EVENT, EXCEED THE ORIGINAL PURCHASE PRICE THEREOF. ECOSCIENCE PRODUCE SYSTEMS CORP. SHALL UNDER NO CIRCUMSTANCES BE LIABLE FOR INCIDENTAL, REMOTE OR CONSEQUENTIAL DAMAGES RESULTING FROM ANY DEFECTS OR ALLEGED DEFECTS IN THIS PRODUCT. EXCEPT TO THE EXTENT THAT AN ALLOWANCE FOR INCIDENTAL DAMAGES IS MANDATED BY APPLICABLE LAW, ECOSCIENCE PRODUCE SYSTEMS CORP. SHALL HAVE NO LIABILITY FOR ANY CLAIM RESULTING FROM THE MISUSE OF THIS PRODUCT OR ANY USE OTHER THAN THAT FOR WHICH IT WAS SPECIFICALLY DESIGNED. No agent of EcoScience Produce Systems Corp. is authorized to make any warranties beyond those contained herein.

EcoScience

PRODUCE SYSTEMS DIVISION

153 Sabal Palm Drive

Longwood, FL 32779

Telephone: 877-866-5773

Facsimile: 407-872-2261

Bio-Save 10 LP

ACTIVE INGREDIENT:

Pseudomonas syringae Strain ESC-10 29.8%

INERT INGREDIENTS: 70.2%

Total 100.0%

Note: Contains a minimum of 9×10^{10} colony forming units per gram of formulated product.

Bio-Save® 10 LP is a naturally occurring biological control agent for postharvest applications only. Do not add directly to mixes, soaps or sanitizers. Do not add to chlorinated water. Application of most chemical fungicides should occur after Bio-Save® 10 LP has been applied. Contact your EcoScience technical advisor for more information.

CITRUS FRUIT (Lemons, Oranges, Grapefruit)

Bio-Save® 10 LP is recommended to aid in the control of green mold (*Penicillium digitatum*), blue mold (*Penicillium italicum*) and sour rot (*Geotrichum candidum*).

Non-recovery Spray: Add 150 grams of product to 10 gallons of water. Agitate the mixture to ensure proper suspension. Apply by drip or spray system to freshly cleaned fruit, prior to waxing. Apply over soft, clean brushes or donut rolls.

CHERRIES

Bio-Save® 10 LP is recommended to aid in the control of blue mold (*Penicillium expansum*), gray mold (*Botrytis cinerea*).

Conventional Dip or Drench: Add 150 grams of product to 10 gallons of water. Agitate the mixture to ensure proper suspension. Drench fruit thoroughly. Recycled dip/drench suspension will need to be recharged at intervals dependent on individual customer use; consult an EcoScience technical advisor for more information.

Overhead Application System: Add 150 grams of product to 10 gallons of water. Agitate the mixture to ensure proper suspension. Apply over conveyor belt or rollers by drip or spray to cherries prior to packaging. Uniform coverage is required. Recycled suspension will need to be recharged at intervals dependent on individual customer use; consult an EcoScience technical advisor for more information. Best control is obtained with an application rate of 1 gallon of suspension to 2,000-4,000 lbs. of cherries.

EPA Reg. No. 68182-xx

EPA Establishment No. 68182

Net Contents: 150 grams

*The biocontrol
Bio-Save is
registered for
postharvest use*

Spectrum of Activity of Biocontrols for Postharvest Decay Control

Biocontrol	Organism	Crops	Decays
Bacteria	<i>Pseudomonas syringae</i>	Apples, pears, citrus	Penicillium Decays
		Sweet cherry	Gray mold, Penicillium decays
Yeast	<i>Candida oleophila</i>	Pome fruit	Penicillium Decays
		Citrus	Penicillium Decays

Biocontrol products registered in other countries

- YieldPlus (*Cryptococcus albidus*) – developed in South Africa for pome fruit
- Avogreen (*Bacillus subtilis*) – South Africa for avocado
- Shemer (*Metschnikowia fructicola*) – Israel for apricot, peach, citrus, grapes, pepper, strawberry, sweet potato
- Several other products such as **Candifruit** (*Candida sake*), **NEXY** (*Candida oleophila*), and **Boni-Protect** (*Aureobasidium pullulans*) are in development.

Postharvest treatments approved for organic produce and their limitations

- | | |
|-----------------------------------------------------------------------------------|-----------------------------------------|
| • Sodium bicarbonate | - Short-lived |
| • Calcium chloride and other chlorine products (with their rates defined by OMRI) | - Only water and surface-disinfestation |
| • Diluted ethanol (not in the US) | - Highly regulated by government |
| • Heat | - Cost, damaging to some crops |
| • UV irradiation | - Cost, damaging to some crops |
| • Biocontrol agents | - Inconsistent |

Prevention, suppression, and eradication of postharvest decays

— Fungicides vs. biological controls —

Fungicides	Biological controls
Single synthetic active ingredient	Mixtures of active and inactive ingredients. Active ingredient often unknown.
Well characterized chemically and toxicologically	Chemically and toxicologically often poorly characterized, but considered natural.
Efficacy generally high	Efficacy variable

Development of Fungicides for Management of Plant Diseases

Initially, developed as simple elements or organic compounds that are non-systemic in plant tissue, and have a low-resistance potential to target organisms.....



but over time, they have been developed as more complex organic compounds, that may be systemic in plant tissue, and have a high-resistance potential to target organisms.

Fungicides have a specific spectrum of activity and, in most cases, are suitable for a limited number of crops

Classes of postharvest fungicides

- Compounds within each fungicide class have:
 - Similar chemical structures
 - A similar mode of action that targets either a single site or multiple sites in the biochemical pathways of the fungus
- Cross-resistance may occur among compounds within the same chemical class

Important older postharvest fungicides for citrus and pome fruits that are still being used today

Residual Fungicide	Class/Grouping	Crops	Decays
SOPP	Phenol	Citrus	Penicillium decay, sour rot
Thiabendazole	Benzimidazole	Citrus, pome fruit	Penicillium decay, gray mold
Imazalil	SBI-Imidazole	Citrus	Penicillium Decays

Towards safer postharvest decay control materials


Re-registration requirements of older pesticides

Reduced Risk Pesticides (an EPA Classification)

- A relative term that is applied to a pesticide as compared to currently registered pesticides of a crop group.
- A pesticide that broadens the adoption of IPM practices or reduces:
 - Exposure risk to humans
 - Potential toxicity to non-target organisms
 - Contamination of the environment

Primarily reduced-risk fungicides will be registered for postharvest use in the US

Benefits of postharvest reduced-risk fungicides to prevent decay



Untreated and postharvest treated (Scholar) peaches and sweet cherries

Spectrum of Activity of Registered and New Postharvest Fungicides on Selected Agricultural Crops in the US

Fungicide	Class	Crops	Decays
Tebuconazole	SBI-Triazole	Sweet cherry	Brown rot, Rhizopus, and Mucor decays
Fludioxonil	Phenylpyrrole	Stone fruit*, pome fruit* Pomegran.*, kiwifruit* citrus, Pineapple, tuber crops	Brown rot, gray mold, Rhizopus Rot, Penicillium decays
Azoxystrobin	Qol	Citrus potato	Penicillium decays
Fenhexamid	Hydroxyanilide	Stone fruit, pome fruit, pomegranate, kiwifruit	Brown rot, gray mold
Pyrimethanil	Anilinopyrimidine	Stone fruit , pome fruit, citrus	Penicillium decays, brown rot, gray mold
Difenoconazole	SBI-Triazole	Pome fruit, tuber crops	Penicillium decays, Bull's eye rot Rhizopus rot
Propiconazole	SBI-Triazole	Stone fruit, citrus, tomato, pepper	Penicillium decays, brown rot, gray mold, sour rot

■ Fungicide is already registered; * - FAT approved in Japan.

new registrations or proposals are in bold italics

Preventing fungicide resistance in the postharvest environment

Guidelines in postharvest fungicide registrations:

Pre-mixtures or tank mixtures of different classes

With mixtures, the resistance potential is much reduced:

$$\begin{array}{|c|} \hline \text{Res. frequency} \\ \text{compound A} \\ \hline \end{array} \times \begin{array}{|c|} \hline \text{Res. frequency} \\ \text{compound B} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{Res. frequency} \\ \text{Pre-mix AB} \\ \hline \end{array}$$

Example:

$$10^6 \times 10^9 = 10^{15}$$

Mixtures of two or three active ingredients that belong to different chemical classes are critical in the prevention of fungicide resistance in target populations.

Postharvest fungicide pre-mixtures

DMI

Imazalil

+

Anilinopyrimidine

Pyrimethanil

=

Philabuster

Citrus

Phenylpyrrole

Fludioxonil

+

QoI

Azoxystrobin

=

Graduate A+

Citrus

Phenylpyrrole

Fludioxonil

+

DMI

Propiconazole

=

Chairman

Stone Fruit, Tomato,

Pineapple

Phenylpyrrole

Fludioxonil

+

MBC

TBZ

=

Scholar Max MP

Pome fruit - registered

Phenylpyrrole

Fludioxonil

+

DMI

Difenoconazole

=

Stadium

Potato

+

Azoxystrobin

and others

Application of postharvest fungicide treatments

- Drenches
- High volume sprayers
- Low volume sprayers (CDA)

Less common:

- Dips
- Flooders
- Foamers
- Brushes
- Fumigators
- Dusters
- Paper wraps
- Box liners

Application of postharvest fungicide treatments

- High volume applications: 100-200 gal/ton of fruit
- Low volume applications: 8-30 gal/ton of fruit

Low volume application systems have become more popular because of very little run-off and no disposal problems

Application methods for postharvest fungicide treatments

High-volume spray application ('T-Jet')



Application methods for postharvest fungicide treatments



Low-volume spray application
(Controlled droplet application - CDA)

Application methods for postharvest fungicide treatments

Dip application



Application methods for postharvest fungicide treatments

Flooder application



Application methods for postharvest fungicide treatments



Flooder
application

Application methods for postharvest fungicide treatments

Fogging



Application of postharvest fungicide treatments

- Aqueous applications
- Application in wax-oil emulsions
 - Not all fruit coatings are considered food-grade in different international markets
 - Prevention of water loss while still permitting gas exchange
 - Increase of shine of fruit

Common fruit coatings used in postharvest treatments

Type of wax	Characteristics			Use on specific crops			
	Prevention of water loss	Gas exchange	Shine of fruit*	Citrus	Nectar./ Peach/ cherry	Plum	Pome
Mineral oil non-emulsified	+++	+	+++		+	+	
Mineral oil emulsified	++	++	+++		+	+	
Polyethylene	+++	+++	+++	+			
Vegetable oils	++	++	++		+	+	
Carnauba	+++	+++	++	+	+	+	+
Shellac	+	+/-	+++	+			+
Wood rosin blends	+	+/-	+++	+			

- Shine of fruit is not important for peaches and plums.
- Carnauba coatings are made from leaves of the Brazilian life tree. Shellac coatings are made from insect exudates. Wood rosins (ester derivatives) are extracted from pine trees.
- Mixtures of polyethylene, carnauba, shellac, and wood rosins are also used on citrus.
- Mixtures of carnauba and shellac are also used on pome fruits.

Postharvest fungicide treatments as a component of postharvest handling

Example: Lemons in California

Fruit arrival



→ Sorting



↓
Chlorine wash,
soda ash
treatment, water
rinse



↓
Application of
fungicide and
fruit coating



Storage wax application



↓

Bulk
packing in
bins

↓


Storage
for up to 3
months



Pack wax application

↓

Boxing, shipping,
marketing





Chlorine wash after storage



Sorting ↓



Boxing and marketing



Fungicide and pack wax application



Use limits of pesticides

Residue tolerance: Maximum residue limit or MRL of a chemical that is allowed on a specific commodity.

Risk assessment based on:

- Toxicological characteristics of chemical
- Amount of human consumption of a specific commodity.

Note - Actual chemical residues are fractions of the tolerances or MRLs

Graduate (fludioxonil) MRLs in major export markets:

	Lemon	Orange	Grapefruit	Tangerine
US	10	10	10	10
CODEX	10	10	10	10
EU	7	7	10	7
Japan FAT	10	10	10	10
Korea	5	5	10	1
Australia	10	10	10	10
Taiwan	7	5	5	—
Following CODEX:	Hong Kong	Hong Kong	Hong Kong	Hong Kong
	India	India	India	India
	Malaysia	Malaysia	Malaysia	Malaysia
	New Zealand	New Zealand	New Zealand	New Zealand
	Philippines	Philippines	Philippines	Philippines
	Singapore	Singapore	Singapore	Singapore
	Thailand	Thailand	Thailand	Thailand

Calculations and Verification for Proper Delivery of Fungicide to Fruit

- Fruit Weight
 - Bin count per time
 - Fruit weight per treatment bed per time
- Fungicide Weight per Volume (Delivery rate)
 - Concentration and Flow Rate
 - * Tank Mix
 - * In-Line Injection
- Sampling and residue measurements of the fungicide on the commodity are ***routinely*** done and ***monitored*** by regulatory agencies

Stewardship of Postharvest Fungicide Treatments

Proper use to ensure food and environmental safety, as well as high-quality nutritious fruits and vegetables.

Prevention of resistance in pathogen populations to fungicides

- *Rotate* between fungicide classes
- *Use* labeled rates
- *Limit* the total number of applications
- *Education* of spectrum of activity
- *Sanitation* is essential in an integrated management program

Conclusions

Chemical treatments in postharvest decay management

- Maximum efforts have been placed on:
 - Food safety (EPA's 'Reduced-risk' fungicides)
 - No mammalian activity at registered rates
 - Lower rates (parts per million quantities)
 - Specific to target plant pathogens
 - Delivery of high quality nutritious fruits and vegetables with minimal losses to growers, packers, and distributors
- Development and proper stewardship of *integrated management programs* cooperatively with land grant research and extension programs and federal/state regulatory agencies.

Use limits of pesticides

- **Residue tolerances** must be established for all postharvest chemical treatments except for those that are exempt:
 - EPA – **Exempt** designation or
 - FDA – **GRAS** (Generally Regarded as Safe) designation

Examples for **GRAS** compounds: chlorine, potassium sorbate, potassium bisulfite, sulfur
- **Residue tolerances - Maximum residue limits (MRLs)**
= The highest amount of a chemical that is allowed to remain on the fruit – determined by EPA.
 - Set below the amount that could pose a health concern.
 - Different for different countries – based on consumer habits and risk analysis
- **Food Additive Tolerances (FATs) – Classification as an ingredient for food use** (country specific, e.g., Japan)

Examples of maximum residue limits (MRLs) - US

Fungicide	MRL	LD ₅₀ rat
Fludioxonil	Stone fruit: 5 mg/kg	>5000 mg/kg
Fenhexamid	Stone fruit: 10 mg/kg	>2000 mg/kg
Pyrimethanil	Citrus: 7 mg/kg	>5000 mg/kg

mg/kg = ppm


Remember that these are maximum levels and actual residue levels are just fractions of these values to obtain desired control.

How many apples does someone need to eat to reach the LD₅₀ of fludioxonil (>5000 mg/kg)?

1 ppm = 1 mg/kg
or 1 mg/10 apples



X 5000 =



50,000 apples/Kg x Body weight (70 kg for an adult)=3.5 million apples!

If you are still concerned....

- **Wash your fruit!**
(Most fungicides are not systemic and can be removed with a household rinse)



Useful Publications - Books:

Postharvest Technology of Horticultural Crops

3rd Edition ANR Publication No. 3311. 2002

Edited by A. A. Kader

Postharvest Pathology

1st Edition Springer, New York, 2010

Edited by D. Prusky and M. L. Gullino

Postharvest: An Introduction to the Physiology and Handling of Fruit and Vegetables

Wills et al., AVI Publishing Co., 1981

A Colour Atlas of Post-harvest Diseases & Disorders of Fruits and Vegetables

A. L. Snowdon, Wolfe Scientific, 1990

Useful Websites (for fungicides):

Labels and MSDS information:

<http://www.cdms.net/manuf/manuf.asp>

<http://www.agrian.com/labelcenter/results.cfm>

Maximum Residue Limit (MRL) or Tolerance information:

<http://www.mrldatabase.com/>

http://ec.europa.eu/sanco_pesticides/public/index.cfm

EPA Fact sheets on new active ingredients:

<http://www.epa.gov/opprd001/factsheets/>

<http://www.epa.gov/oppfead1/trac/safero.htm>

Research:

<http://californiaagriculture.ucanr.org/Landingpage.cfm?article=ca.v059n02p109&fulltext=yes>

Useful Websites (Postharvest Companies):

Service companies -

Decco:

<http://www.deccous.com/>

JBT (formerly FMC):

<http://www.jbtfoodtech.com/solutions/equipment/fresh-produce-technologies/post-harvest-products-and-services.aspx>

Pace International:

<http://www.paceint.com/>

Fungicide companies -

Syngenta Postharvest University:

<http://www.farmassist.com/postharvest/index.asp?nav=contact>

Janssen PMP:

<http://www.janssenpmp.com/>